



ZIBELINE INTERNATIONAL

Print ISSN : 2521-0858

Online ISSN : 2521-0866

CODEN: SHJCAS



THE POTENTIAL OF DRIED AND FRESH EXTRACTS OF SANSEVIERIA TRIFASCIATA TO MITIGATE ALEXANDRIUM TAMIYAVANICHII, A TOXIC DINOFLAGELLATE

Normawaty Mohammad Noor^{1*}, Ima Amirah Mohd Suberi¹, Deny Susanti², Yukinori Mukail¹, Aimimuliani Adam¹, Shahbudin Saad¹, Fikri Akmal Khodzori¹

¹Department of Marine Science, Kulliyah of Science International Islamic University Malaysia, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia ·

²Department of Chemistry, Kulliyah of Science International Islamic University Malaysia, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia.

*Corresponding Author's Email: normahwaty@iiu.edu.my

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

ARTICLE DETAILS

Article History:

Received 12 November 2017

Accepted 12 December 2017

Available online 1 January 2018

ABSTRACT

Dinoflagellate is the second largest group of phytoplankton in the marine environment and many of the species have been recognized as toxic species. The high abundance of this species can cause a lot of problems to human health, marine organisms especially caged fish and to the natural environment. In Malaysia, several species of *Alexandrium* have been reported to cause human intoxication after consumption of contaminated shellfish. During the event, shellfish was banned and lead to economic lost especially to the local fishermen. Therefore, this study was conducted to find alternative way to mitigate the blooms of harmful algae particularly *Alexandrium tamiyavanichii* using ornamental plant, *Sansevieria trifasciata* or known as mother-in-law tongue. The extraction of this plant was done in distilled water using dried and fresh plants. Different concentrations of crude extracts (0.001, 0.01, 0.1, 0.5, 1, 1.5, 2 and 2.5 mg/mL) were tested on *A. tamiyavanichii* within 24 hours. The removal efficiencies (RE) were determined for each concentration tested. The extracts were also tested on brine shrimp to see the possible harmful effect. Results indicate that both extraction methods yielded positive results on *A. tamiyavanichii* whereby the crude extracts managed to mitigate the species. The LC50 of brine shrimp recorded were 30 mg/ml for fresh plant extract and 70 mg/mL for dried plant extracts. This indicates the possibility of the extractions to be used to mitigate harmful algal bloom (HAB) particularly *Alexandrium* and provide an alternative way to the relevant agencies to minimize the impact of HAB

KEYWORDS

Alexandrium tamiyavanichii, *Sansevieria trifasciata*, distilled water crude extract, removal efficiency.

1. INTRODUCTION

Harmful algal bloom (HAB) is an aquatic phenomena that occur when there is discoloration of water caused by bloom of microscopic algae whether in marine or freshwater environment. Occurrence of the bloom can cause enormous lost in aquaculture industries, economic and also harmful to human and animal health. For decade, this problem is globally occurred. In Malaysia, several cases have been reported relate to contaminated shellfish by toxic algae species involving *Alexandrium* [1]. Many studies have been conducted to find the best method for HABs mitigation that is feasible, cost-effective and has minimum impacts to the environment [2]. Most popular method is applying clay to the infected area. This method is very famous and has been applied in several countries such as Korea, Japan, China and Thailand [3]. Other than that, biological approaches like algicidal bacteria (*Micrococcus luteus*) also have been used to reduce the damaged caused by harmful algae [4]. All of these methods produced positive results and some showed specific effect on certain species [5]. However, some of the methods mentioned will cause pollution to the environment and affect others marine life. Nowadays, the use of allelopathy for HABs control has been given attention due to its low cost and environmental friendly [6].

In Malaysia, study on HABs control still at initial stage. Looking at the impact of HAB on human health and economy [1][7], it is an urgent need to identify potential way to mitigate HAB species. An ornamental plant which is *Sansevieria trifasciata* or known as snake plant was chosen for the study as the preliminary study on mitigation of HAB showed a positive result [unpublished data]. This plant is popular in traditional medicine and being used in many studies due to its benefits in medicinal, fodder, soil conservation and fibre [8].

Therefore, in this study, crude extract of *S. trifasciata* was used to mitigate a toxic dinoflagellate, *A. tamiyavanichii* isolated from Kuantan, Pahang. Besides, the LC50 of the crude extracts on brine shrimp were determined to represent its effect on other marine organisms. Through this study, an alternative way for HABs mitigation particularly *A. tamiyavanichii* can be suggested. The outcomes of the study are expected to contribute significantly on mitigation of HAB particularly in Malaysia.

2. EXPERIMENTAL

2.1 *A. tamiyavanichii* culture

Cells of *A. tamiyavanichii* were isolated from samples collected from Kuantan Port. The cultures were grown at 26°C with 12:12 L: D (light and dark) cycle and maintained in ES-DK medium enriched with f/2 vitamin. The light intensity used was 1000 lux.

2.2 Extraction of *S. trifasciata*

Plants were washed, weighted and cut. To obtain dried sample, plants were dried in the oven at 40°C until constant weight were obtained. Then, both samples (fresh and dried) were grinded using mixer before soaked in distilled water and heated in water bath at 60°C for 8 hours. Then, crude extracts were filtered before placed in -80°C freezer and finally undergo freeze drying.

2.3 Test of extracts on *A. tamiyavanichii*

These crude extracts were tested on *A. tamiyavanichii* at different concentrations (0.001, 0.01, 0.1, 0.5, 1, 1.5, 2.0 and 2.5 mg/mL). The effects of each concentration of the crude extracts were determined at time interval of 0, 2.5, 5, 10 and 24 hours. The cell density tested was approximately 1000 to 2000 cells/mL. Tests were conducted in 60×10 mm Petri dish with 3 replicates and 1 control for each concentration. pH of the medium after mixed with the crude extracts were measured for each concentration. Numbers of cells were counted twice under light microscopy at total magnification of 100× using Sedgewick rafter cell for each concentration and time tested. Finally, the effects of the crude extracts on the growth of *A. tamiyavanichii* were determined based on removal efficiency (RE). The formula used as follows. The higher removal efficiency means that the extract is more effective in removing the cells. $RE = [(initial\ cell\ concentration - sample\ cell\ concentration) / initial\ cell\ concentration] \times 100\%$

2.4 Toxicity test

Brine shrimp usually being used in toxicity test because its lethality bioassay was rapid, easily to get, simple and cheap. The cysts were hatched in sterile seawater for 24 hours before tested with different concentrations of crude extracts which were 10, 50, 100 and 500 mg/mL. The tests were conducted in 60×10 mm petri dish with 3 replicates and 1 control. There were 10 tails of brine shrimp in each Petri dish. LC50 at the 95% confidence limits were obtained by plotted survivors' percentages with concentrations tested.

2.5 Statistical analysis

Results were analysed by one-way ANOVA using SPSS 20.0 to see if there were significant differences between concentrations and times tested for both extracts. Correlation analysis was performed to see the relationship of REs with concentrations and time exposed.

3. RESULTS AND DISCUSSION

3.1 Test of extracts on *A. tamiyavanichii*

Figures 1 and 2 showed the results of removal efficiency of fresh and dried plant extracts on *A. tamiyavanichii*. The highest result of removal efficiency of fresh plant extract on *A. tamiyavanichii* was recorded at concentration of 2 mg/mL which was 88.1% at 0 hours whereas the lowest removal efficiency was 26.85% that tested with 0.001 mg/mL concentration at 2.5 hours (Figure 1). The pHs for both concentrations after mixed the extracts with medium containing *A. tamiyavanichii* were 5.96 and 8.27, respectively (Table 1).

The removal efficiency of dried plant extract on *A. tamiyavanichii* showed that the algae exposed to 1.5 mg/mL concentration had the highest result which was 78.24% at 5 hours while the lowest removal efficiency was 23.28% tested with 0.1 mg/mL plant crude extract at 24 hours (Figure 2). The pHs recorded were 6.27 and 8.28 for concentration 1.5 and 0.1 mg/mL after mixed the extract with media containing *A. tamiyavanichii* (Table 1). Both extracts showed significant differences ($p < 0.05$) of removal efficiency between times and concentrations for all experiment conducted.

Results showed that *A. tamiyavanichii* reacts differently on different concentrations of both extracts tested. Statistical analysis showed a positive correlation ($r = 0.76, r = 0.67$) between removal efficiency and concentration for both crude extracts tested. However, between removal efficiency and time, negative correlations ($r = -0.29, r = -0.44$) were obtained. Previous study on harmful algae, *P. globosa* and *P. donghaiense* using Chinese traditional herbs and herb-modified clay have showed that the inhibitory rate of algae increased as the concentration applied increased [9].

Result obtained also showed that as the concentration increased, pH decreased and removal efficiency increased (Table 1). The dropped of medium pH after the extracts were added need to be given attention. This is because different species has different tolerant towards pH [10]. This is to ensure that high RE recorded is due to the effect of crude extracts tested and not change in pH of the medium.

Table 1: Result summary of plant extracts

	Fresh plant extract		Dried plant extract	
	highest	lowest	highest	lowest
Concentration (mg/mL)	2	0.001	1.5	0.1
RE (%)	88.1	26.85	78.24	23.28
Time (hour)	0	2.5	5	24
pH	5.96	8.27	6.27	8.28
LC50	30 mg/mL		70 mg/mL	

Both extracts showed significant differences ($p < 0.05$) of removal efficiency between times and concentrations for all experiment conducted. Results showed that *A. tamiyavanichii* reacts differently on different concentrations of both extracts tested. Statistical analysis showed a positive correlation ($r = 0.76, r = 0.67$)




Figure 1: Removal efficiencies (RE) (%) of fresh plant extracts on *A. tamiyavanichii*




Figure 2: Removal efficiencies (RE) (%) of dried plant extracts on *A. tamiyavanichii*

3.2 Toxicity test

Figure 3 showed the results of LC50 of distilled water fresh and dried plant extracts on brine shrimp (artemia). The LC50 of distilled water fresh plant extract on artemia was lower i.e. 30mg/mL compared to dried plant extract i.e. 70 mg/mL (Figure 3). From the toxicity result, high concentrations of extracts were needed to kill the brine shrimp. This study help to prove that the extracts might be suitable to be applied at the field as the concentration needed to kill harmful algae is very low compared to the LC50. Nevertheless, further study on other organisms such as to bivalve and farmed fish need to be conducted to ensure that the extracts have no effect on surrounding flora and fauna [11].




Figure 3: LC50 of fresh plant extract (A) and dried plant extract (B) on artemia

4. CONCLUSION

Results of removal efficiencies showed different effectiveness between both crude extracts (fresh and dried plant extract of *S. trifasciata*) on *A. tamiyavanichii*. However, overall results showed that the pH influenced the results by increasing the removal efficiency. Therefore, high concentration of crude extracts is not recommended especially for toxic algae. This is because the cells will lyse and toxin will be released to the environment posing other secondary effect. Nevertheless, this study indicates that ornamental plant, *S. trifasciata* has the potential to inhibit the growth of *A. tamiyavanichii* but further study is needed to fine tune the result in order to find the suitable concentration needed to mitigate HAB species. Results of this study are limited to *Alexandrium tamiyavanichii* and based on 2 concentrations tested.

ACKNOWLEDGEMENT

We would like to thank Kuliyyah of Science, International Islamic University Malaysia for all the facilities provided.

REFERENCES

- [1] Lim, P.T., Usup, G., Leaw, C.P. 2012. Harmful algal bloom in Malaysia waters, Sains Malaysiana, 41, 1509-1515.
- [2] Sengco, M.R., Anderson, D.M. 2004. Controlling harmful algal blooms through clay flocculation. Journal of Eukaryotic Microbiology, 51, 169-172. DOI: 10.1111/j.1550-7408.2004.tb00541.x.
- [3] Pan, G., Yang, B., Wang, D., Chen, H., Tian, B.H., M. L. Zhang, M.L., Yuan, X.Z., Chen, J. 2011. In-lake algal bloom removal and submerged vegetation restoration using modified local soils. Ecological Engineering, 37, 302-308. DOI: 10.1016/j.ecoleng.2010.11.019. 10.1007/s10811-013-0046-z.
- [4] Min-Ju, K., Seong-Yun, J., Sang-Joon, L. 2008. Isolation, identification and algicidal activity of marine bacteria against *Cochlodinium polykrikoides*. Journal of Applied Pshycology, 20, 1069-1078. DOI: 10.1007/s10811-008-9312-x.
- [5] Baek, S.H., Son, M., Jung, S.W., Na, D.H., Cho, H., Yamaguchi, M., Kim, S.W., Kim, Y.O. 2013. Enhanced species-specific chemical control of harmful and non-harmful algal bloom species by thiazolidinedione

derivative TD49. Journal of Applied Pshycology, 26, 311-321

[6] Renjun, W., Xiao, H., Zhang, P., Qu, L., Cai, H., Tang, X. 2007. Allelopathic effects of *Ulva pertusa*, *Corallina pilulifera* and *Sargassum thunbergii* on the growth of the dinoflagellates *Heterosigma akashiwo* and *Alexandrium tamarense*. Journal of Applied Pshycology. 19, 109-121 DOI: 10.1007/s10811-006-9117-8.

[7] Adam, A., Mohammad-Noor, N., Anton, A., Saleh, E., Saad, S., M. Shaleh, S.R. 2011. Temporal and spatial distribution of harmful algal bloom (HAB) species in coastal waters of Kota Kinabalu, Sabah, Malaysia. Harmful Algae, 10 (2011), 495-502. DOI: 10.1016/j.hal.2011.03.006.

[8] Khalumba, M.L., Mbugua, P.K., Kung'u, J.B. 2005. Uses and conservation of some highland species of the genus *Sansevieria* Thunb in Kenya. In African Crop Science Conference Proceeding, 7, 527-532.

[9] Tian, F., Zhou, J., Sun, Z., Cai, Z., Xu, N., An, M., Duan, S. 2014. Inhibitory effects of Chinese traditional herbs and herb-modified clays on the growth of harmful algae, *Phaeocystis globosa* and *Prorocentrum donghaiense*, Harmful algae, 37, 153-159. DOI: 10.1016/j.hal.2014.05.015.

[10] Berge, T., Daugbjerg, N., Andersen, B.B., Hansen, P.J. 2010. Effect of lowered pH on marine Phytoplankton growth rates. Marine Ecology Progress Series, 416, 79-91. DOI: 10.3354/meps08780.

[11] Z. Yu, X. Song, X. Cao, Y. Liu, Mitigation of harmful algal blooms using modified clays: theory, mechanism and application. Harmful Algae. 69 (2017), 48-64.

