

REVIEW ARTICLE

ALGAL DIVERSITY IN NEPAL AND ITS APPLICATIONS: CURRENT INSIGHTS AND FUTURE PROSPECTS

Sandesh Subedi

*Institute of Agriculture and Animal Science, Lamjung Campus (Tribhuvan University), Lamjung, Nepal.***Corresponding Author Email: sandesh.181401@lac.tu.edu.np*

This is an open access article distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 20 December 2023

Revised 15 January 2024

Accepted 02 February 2024

Available online 05 February 2024

ABSTRACT

Algae are a diverse group of unicellular autotrophic microorganisms. It is a free-floating and filament-shaped organism, in which the blade is the main site for photosynthesis. Hence, called a photosynthetic organism. Different types of algae are found in different types of climatic conditions. Algae are found in almost all adverse climates. Due to its vast chemical composition and content of bioactive substances attention of different industries like food, cosmetic pharmaceuticals, nutraceuticals, etc. are caught by it. Algae food products contain a high amount of nutrients, and cosmetic products protect from UV radiation and prevention of rough texture wrinkles of the skin, as they carry out both photosynthesis as well as nitrogen fixation, add organic matter (20 -30kg/ha/year) and maintain texture porosity, the structure of soil and can play an important role for reducing the problem of petrol, diesel. This review has been prepared to shed light on the wide uses of algae as food, fuel, cosmetic product biofertilizer and also to provide knowledge about different types of algae found in different geographical zones of Nepal.

KEYWORDS

Algae, biofuel, seaweeds, unicellular

1. INTRODUCTION

Linnaeus first derived Algae from the Latin word "Alga" which means sea weeds. The study of algae in scientific terms is called Phycology/Algology (Lee, 2008). It is commonly called Children of the Sea. F.E. Fritsch is considered the father of Phycology. Chlorella is the smallest algae and Acetabularia (sea umbrella) is the largest unicellular algae. Poryphyra linearis has the smallest number of chromosomes i.e. $n=2$, whereas the highest number of chromosomes is found in Natrium digitalis i.e. $n=592$. Algae are green, aquatic chlorophyll-containing photosynthetic organisms but lacks true stem, and leaves, and roots. About 90% of the total photosynthesis is performed by algae. Algae are simple plants with thallus structures. They are considered as most primitive organisms that absorb toxic substances.

Algae are chlorophyllous and non-vascular. It is found in every type of soil, on and within other plants, even animals, rocks, on stones, and on diverse climates where a large amount of water is available i.e. both marine and freshwater (Korbitz et al., 1994). In algae, the embryo stage is absent, and non-jacketed gametangia (gametes) are present where all the cells are fertile and meiosis occurs by a zygotic method. They are unicellular, colonial, filamentous, motile, non-motile, dendroid, coccoid, heterotrichous etc. Most of the algae are microscopic, and unicellular and their maximum diameter ranges up to 0.5μ and 60m in length. Algae are covered by mucilages similar to other aquatic plants and protects the plants through decaying of water (Black and McGinnis, 2006). Moreover, it also protects from epiphytic growth and desiccation (Elnabris, 2000). The cell wall made up of pectin, chitin, algin, and fucondin surrounds each cell. Vacuoles, chloroplast, nucleus, pyrenoids, Golgi bodies, and various types of pigments are embedded inside the cell wall. Starch is the reserved food material in most of the algae, which also varies according to different forms. Reproduction in algae occurs by sexual, asexual, and vegetative methods. A vegetative method includes fragmentation, fission, akinete, tuber, and hormogonia. Asexual reproduction involves the division of

protoplast and this protoplast grows into new plants. It occurs by forming spores like zoospores, syn-zoospores, aplnospores, autospores, auxospores, etc. Sexual reproduction is the fusion of male and female gamete. It is of isogamous, anisogamous, and oogamous (Bhuju et al., 2065).

Algae are unicellular microscopic organisms i.e. microalgae and can extend to greater size i.e. macro algae (Munir et al., 2013). Microalgae consists of diatoms, some green algae, dinoflagellates, euglenids, and some red algae and contain single flagella. They are free-floating and their body shape appears to be filaments. Some microalgae like volvox live in a colony. Macro algae consist of brown algae, green algae, and red algae. It contains holdfast which helps to keep the plant in one place (Fall et al., 2010). Blades are the main sites of photosynthesis in algae (Roberts et al., 2007). Algae are found in both land and water environments. Epiphyllaphytes, epiphyllaphytes, and lithophytes, are the terrestrial form of algae whereas phytoplanktons, benthic/ Cryptophytic, epiphytic, epizotic, endozoic, and periphytic are the different aquatic algae. Cryophytes are those algae which can sustain in low temperature i.e. on snow E.g. *Haematococcus*, *Scotiella* etc whereas thermophytes can tolerate high temperature i.e. on hot spring water E.g. *Oscillatoria brevis* (Blue green algae), *Haplosiphon*.

Algae plays a major role in the purification of the environment, to meet the current demand for biofuels and reduce the pressure of petroleum and diesel fuels, which are going to be exhausted soon. More over when food is scarce, algae can provide the nutrient diets to human beings and animals. Algae reduces the concentration of carbon dioxide in the atmosphere and gives oxygen which is one of the most essential requirements for the existence of life. Algae act as biofertilizers and improve the soil structure, porosity, and texture. Water blooms produced by the Blue Green Algae and Green algae lead to the death of aquatic organisms (Hasan and Chakrabarti, 2009). Red rust of tea and coffee is caused due to *Cephaleuros virescens* and *Cephaleuros arabica*.

Quick Response Code



Access this article online

Website:

www.jscienceheritage.com

DOI:

[10.26480/gws.02.2024.69.78](http://doi.org/10.26480/gws.02.2024.69.78)

1.1 Types Of Algae

Although alga looks simple in forms and structures they have diverse classifications. According to 11 classes are present based on pigment and reserve food materials of algae, which are as follows (Fritsch, 1935)

Table 1: Classification of algae (Elnabris, 2000)										
Name	Occurrence	Orders	Structure	Pigments	Reserve food	Pyrenoids	Cell wall	Flagella	Reproduction	Examples
Eugleuineae	Fresh water	-	Unicellular	Chl-a,Chl-c,β carotene	Polysaccharide and paramylon	Similar structures	Proteinaceous	One or two	Cell division, Sexual rare(isogamous)	<i>Euglena</i>
Chloromonado phyceae	Fresh water	1	Motile unicells	Chl-a, xanthophylls	Fats or oils	Absents	Absent	Biflagellate equal	Cell division, Sexual reproduction absent	<i>Gonyostomum</i>
Bacillariophyc eae	Cosmopolitan distribution, even in soil, terrestrial	2	Unicellular or colonial	Chl-a,Chl-c,β carotene Fucoxanthin, Diatoxanthin, didinoxanthin	Fats, Volutin	Present	Siliceous(outer) and pectic(inner)	Single flagella	Cell division and auxospore	<i>Pinnularia</i> , <i>Cymbella</i> , <i>Navicula</i>
Cryptophyceae	Both freshwater and marine	2	Unicellular with anterior groove or pocket	Chl-a,Chl-c, Diatoxanthin	Starch or oil	Pyrenoid like structures is present.	Absent	Biflagellate, equal or unequal	Binary fission, sexual rare(isogamous)	<i>Chroomonas</i> , <i>Cryptomonas</i> ,
Chrysophyceae	Mostly freshwater, few marine	3	Unicelled motile to filamentous branched	Chl-a,Chl-c,β carotene, Phycocytin	Leucosin, fats,Chrysolaminarin	Absent	Siliceous or calcified, cellulose absent	2 equal or unequal flagella	Vegetative,asexual, sexual rare(isogamous)	<i>Prymnesium</i> , <i>Dinobryon</i> , <i>Chrysosphaera</i>
Xanthophyceae	Mostly freshwater, few marine	4	Unicelled motile to filamentous	Chl-a, Chl-e and β carotene, xanthophyll, neoxanthin, and violaxanthin	Oil	Absent	Rich in Pectin	2 unequal flagella (heterokont)	Vegetative, asexual, sexual, (mainly isogamous, anisogamous rare,oogamy in <i>Vaucheria</i>)	<i>Chlorogloea</i> , <i>Tribonema</i> , <i>Chlorogloea</i> , <i>Botrydium</i>
Myxophyceae	Fresh water	5	Unicellular, prokaryotic	Chl-a, xanthophylls phycoerythrin, and blue phycocyanin, carotenes	Cyanophycan starch		Peptidoglycan	Absent	Asexual reproduction by fragmentation, fission, akinetes, hormogonia	<i>Oscillatoria</i> , <i>Nostoc</i>
Dinophyceae	Mostly marine	6	Unicellular, branched filamentous	Chl-a,Chl-c,β carotene, xanthophyll peridinin, neoperidinin etc	Starch or fats		Cellulose	Biflagellate, equal	Sexual rare(isogamous)	<i>Ceratium</i> , <i>Noctiluca</i> , <i>peridinium</i>
Rhodophyceae	Mostly marine, rarely fresh water	7	Unicellular to complex, Parenchymatous to non parenchymatous thallus seaweeds	Chl-a,Chl-d, phycoerythrin and blue phycocyanin, carotenes.	Floridean starch		Outer wall polysaccharides, inner wall cellulose	Absent	Advance oogamous	<i>Thorea</i> , <i>Gelidium</i> , <i>Champia</i> , <i>Ceramium</i> , <i>Corallina</i>
Chlorophyceae	Aquatic, mostly freshwater, few are terrestrial	9	Unicellular, multicellular to hetetrotichus, filamentous	Chl-a,chl-b, and αβcarotenes, lycopene, xanthophyll	Starch	Present	Cellulosic	Motile cells, equal flagella	Vegetative, asexual, sexual(isogamous, anisogamous to advanced oogamous)	<i>Volvox</i> , <i>pediastrum</i> , <i>ulothrix</i> , <i>cladophora</i> , <i>cephaleuros</i> , <i>Chara</i> , <i>Oedogonium</i>
Phaeophyceae	Marine	9	Multicellular to giant sea weeds (kelps)	Chl-a,Chl-c,β carotene, xanthophylls	Mannitol, laminarin and oils		Polysaccharides alginates & fucose-containing sulfated polysaccharides	Motile gametes 2 laterally attached flagella heterokont	Asexual(fragmentation or zoospores) Sexually (isogamous,oogamous or anisogamous)	<i>Ectocarpus</i> , <i>Tilopteris</i> , <i>Cutieria</i> , <i>Sphaecelaria</i> , <i>Sargassum</i> , <i>Dicotyota</i> , <i>Sporochnus</i>

1.2 Diversity of Algae in Nepal

Algae can suit from adverse climate to favourable environment. In all districts, few to many varieties are found. Following are the districts where research activities have been performed-




Figure 1: Distribution of Algae in Nepal

Due to the difficult land structures, climate and technology most of the research activities are performed in city areas which results for the hinderance of algae found in the remote areas. Some efforts have been

made by different researchers to find different types of algae in different geographical regions are mentioned below-

Table 2: Algae diversity along with districts of Nepal

S.N	Location	Name	References
1	Bara	<i>Gloeocapsa decorticans</i> , <i>Gloeotheca rupestris</i> , <i>Aulosira fertilissima</i> , <i>Aulosira fertilissima</i> , <i>Nostoc commune</i>	(Prasad, 1996)
2	Bhaktapur	<i>Schizothrix purpurascens</i> , <i>Nostoc sphericum</i> , <i>Scytonema fertilia</i> , <i>Hapalosiphon intricatus</i> , <i>Stigonema hormoides</i> , <i>Westiellopsis species</i> , <i>Nostoc linckia</i> , <i>Scytonema bewsii</i> , <i>Calothrix fusca</i> , <i>Aulosira implexa</i> , <i>Phormidium autumnale</i> , <i>Cylindrospermum majus</i>	(Shrestha and Manandhar, 1983; Watanabe and Komarek, 1988)
3	Chitwan	<i>Aphanocapsa elachista</i> , <i>Aphanocapsa rivularis</i> , <i>Arthrospira spirulinoides</i> , <i>Lyngbya lattissima</i> , <i>Anabaena sphaerica</i> , <i>Aulosira fritschii</i> , <i>Aulosira prolifica</i> , <i>Aulosira fertilissima</i> , <i>Microcoleus chthonoplastes</i> , <i>Synechocystis aquatilis</i> , <i>Oscillatoria sancta</i> , <i>Oscillatoria granulate</i> , <i>Oscillatoria princeps</i> , <i>Synechocystis pevalekii</i> , <i>Oscillatoria irrigua</i> , <i>Chroococcus minutus</i>	(Das and Verma, 1996)
4	Gorkha	<i>Dactylococcopsis raphidioides</i> , <i>Merismopedia tenuissima</i> , <i>Oscillatoria granulata</i>	(Hirano, 1955)
5	Illam	<i>Microcoleus sociatus</i> , <i>Phormidium subfuscum</i> , <i>Hapalosiphon fontinalis</i> , <i>Lyngbya hieronymusii</i>	(Rai and Misra, 2010)
6	Jhapa	<i>Chroococcus schizodermaticus</i> , <i>Gloeocapsa nigrescens</i> , <i>Tolypothrix distorta</i> , <i>Rivularia minutula</i> , <i>Eucapsis himalayensis</i>	(Rai and Misra, 2010)
7	Kanchanpur	<i>Aphanocapsa roeseana</i> , <i>Aphanothece microscopica</i> , <i>Microcystis marginata</i> , <i>Lyngbya contorta</i> , <i>Oscillatoria formosa</i> , <i>Phormidium incrustatum</i> , <i>Phormidium usterii</i> <i>Aulosira fertilissima</i> , <i>Gloeotrichia natans</i> , <i>Lyngbya hieronymusii</i> , <i>Lyngbya allorgei</i> , <i>Anabaena iyengarii</i> , <i>Nostoc linckia</i> , <i>Nostoc linckia</i> , <i>Lyngbya hieronymusii</i> , <i>Lyngbya allorgei</i> , <i>Anabaena iyengarii</i> .	(Habib, 1997)
8	Kaski	<i>Anabaena circinalis</i> , <i>Microcystis aeruginosa</i> , <i>Microcystis flosaqua</i> , <i>Spirulina gigantea</i>	(Nakanishi, 1986; Hickel, 1997)
9	Kathmandu	<i>Aphanocapsa anodontae</i> , <i>Aphanothece pallida</i> , <i>Gloeocapsa aeruginosa</i> , <i>Gloeocapsa atrata</i> , <i>Gloeocapsa dermochroa</i> , <i>Gloeocapsa sanguinea</i> , <i>Gloeotheca fusco-lutea</i> , <i>Holopedium irregulare</i> <i>Nephrococcus species</i> , <i>Cyanosarcina species</i> , <i>Lyngbya lagerheimii</i> , <i>Lyngbya maior</i> , <i>Microcoleus vaginatus</i> , <i>Oscillatoria acuminata</i> , <i>Oscillatoria granulate</i> , <i>Oscillatoria proteus</i> , <i>Oscillatoria subproboscidea</i> , <i>Phormidium ambiguum</i> , <i>Phormidium fragile</i> , <i>Schizothrix friesii</i> , <i>Schizothrix heufleri</i> , <i>Schizothrix pulvinata</i> , <i>Anabaena fertilissima</i> , <i>Anabaena globosa</i> , <i>Pseudanabaena catenata</i> , <i>Scytonema arcangelii</i> , <i>Scytonema contorta</i> , <i>Scytonema myochrous</i> , <i>Scytonema schmidtii</i> , <i>Scytonema stuposum</i> , <i>Tolypothrix arenophila</i> , <i>Tolypothrix phyllophila</i> , <i>Calothrix fusca</i> , <i>Calothrix scytonemicola</i> , <i>Oscillatoria foreau</i> , <i>Oscillatoria okeni</i> , <i>Nostoc linckia</i> , <i>Scytonema bewsii</i> , <i>Oscillatoria granulate</i> , <i>Microcystis flosaqua</i> , <i>Phormidium anomala</i> , <i>Calothrix fusca</i> , <i>Rivularia minutula</i> , <i>Lyngbya allorgei</i> , <i>Phormidium uncinatum</i> , <i>Oscillatoria anguina</i> , <i>Nodularia harveyana</i> , <i>Anabaena doliolum</i> , <i>Oscillatoria jasorvensis</i> , <i>Phormidium retzii</i> , <i>Cylindrospermum minutissimum</i> , <i>Cylindrospermum stagnale</i> , <i>Nostoc calcicola</i> , <i>Nostoc ellipsosporum</i> , <i>Calothrix parietina</i> , <i>Merismopedia glauca</i> , <i>Chroococcus turgidus</i> , <i>Gomphosphaeria aponina</i> , <i>Microcystis robusta</i> , <i>Aphanocapsa grevillea</i> ,	(Baral et al., 1988; Hirano, 1963; Watanabe and Komarek, 1988; Watanabe, 1995)

		<i>Oscillatoria irrigua, Merismopedia punctate, Oscillatoria proboscidea, Lyngbya aerugineocoerulea, Nostoc spongiaeforme, Oscillatoria limosa, Chroococcus limneticus.</i>	
10	Khotang	<i>Oscillatoria amoena, Oscillatoria curviceps, Phormidium anomala, Oscillatoria amoena, Oscillatoria curviceps, Spirulina major.</i>	(Rai and Misra, 2010)
11	Lalitpur	<i>Aphanocapsa pulchra, Chroococcus minor Gloeocapsa muralis, Chamaesiphon confervicola, Chamaesiphon minutus, Blennothrix ganeshii, Lyngbya bipunctata, Oscillatoria acutissima, Phormidium corium, Nostoc muscorum, Nostoc paludosum, Scytonema bewsii, Scytonema ocellatum, Homoeothrix janthina, Scytonema bewsii, Aulosira implexa, Oscillatoria princeps, Cyndrospermum stagnale, Nostoc calcicola, Nostoc ellipsoforum, Lyngbya birgei, Nostochopsis lobatus, Oscillatoria amoena, Oscillatoria subbrevis, Oscillatoria chlorine, Calothrix parietina, Microcystis stagnalis, Anabaena variabilis.</i>	(Joshi, 1977; Joshi, 1979; Prasad and Prasad, 2001; Rai and Misra, 2010)
12	Lamjung	<i>Lyngbya confervoides, Oscillatoria deflexa, Oscillatoria laete-virens, Phormidium jadinianum, Phormidium stagnina, Anabaena flosaquae, Microcoleus lacustris</i>	(Panta and Gupta, 1999)
13	Makawanpur	<i>Dactylococcopsis acicularis, Oscillatoria annae, Merismopedia elegans, Spirulina meneghiniana</i>	(Sahay et al., 1993)
14	Manang	<i>Aphanocapsa delicatissima, Oscillatoria amphibian, Oscillatoria terebriformis, Phormidium papyraceum, Phormidium purpurascens, Arthronema africanum, Phormidium autumnale, Calothrix parietina, Merismopedia glauca, Chroococcus turgidus, Gomphosphaeria aponina.</i>	(Hirano, 1955; Komarek and Lukavsky, 1988)
15	Morang	<i>Arthrospira khannae, Lyngbya majuscula, Calothrix castelli, Gloeotrichia intermedia, Fischerella epiphytica, Oscillatoria sancta Oscillatoria anguina, Lyngbya birgei, Nostochopsis lobatus, Microcystis robusta, Aphanocapsa grevillea, Spirulina subsalsa, Spirulina princeps.</i>	(Rai and Misra, 2010)
16	Mugu	<i>Aphanothece stagnina, Leptolyngbya perelegans, Leptolyngbya rivulariarum, Anabaena lapponica, Anabaena oscillarioides, Pseudanabaena mucicola, Microchaete tenera, Calothrix gloeocola, Phormidium tenue, Nodularia harveyana, Cyndrospermum minutissimum, Chroococcus minutus, Rivularia hangirgi</i>	(Watanabe, 1995)
17	Mustang	<i>Synechococcus aeruginosus, Phormidium laminnatum, Gomphosphaeria aponina, Synechococcus aeruginosus, Phormidium laminnatum, Oscillatoria irrigua, Merismopedia punctate, Oscillatoria proboscidea, Microcystis stagnalis, Chroococcus minutus.</i>	(Hirano, 1955; Upadhyaya, 1979)
18	Nawalparasi	<i>Oscillatoria borneti, Anabaena augstumalis, Anabaena unispora, Calothrix atricha, Calothrix castelli, Spirulina gigantean, Oscillatoria agardhii, Anabaena variabilis</i>	(Das and Verma, 1996)
19	Parsa	<i>Gloeocapsa rupestris, Arthrospira platensis, Phormidium lucidum, Anabaena azollae, Anabaenopsis raciborskii, Raphidiopsis curvata, Gloeocapsa decorticans, Gloeotheca rupestris</i>	(Prasad, 1996)
20	Rasuwa	<i>Aphanocapsa elachista, Chamaesiphon incrustans, Hydrocoleus homoeotrichum, Lyngbya epiphytica, Lyngbya sordida, Oscillatoria brevis, Oscillatoria pseudominima, Phormidium valderiae, Schizothrix lacustris, Anabaena papillosa, Scytonema mirabile, Aphanocapsa grevillea, Oscillatoria splendida, Lyngbya aerugineocoerulea, Stigonema ocellatum, Stigonema mamillosum, Oscillatoria animalis</i>	(Hirano, 1969; Yoshimura et al., 1997; 2006)
21	Rautahat	<i>Chroococcus limneticus, Eucapsis minuta, Merismopedia minima, Synechococcus Elongates, Arthrospira massartii, Oscillatoria articulate, Hapalosiphon welwitschii, Gloeocapsa decorticans, Gloeotheca rupestris, Merismopedia punctate, Microcoleus subtorrulosus, Nostoc microscopicum, Cyndrospermum majus, Anabaena iyengarii, Chroococcus tenax, Chroococcus minutus</i>	(Sahay et al., 1993)
22	Sankhuwasabha	<i>Scytonema burmanicum, Stigonema ocellatum, Chlorogloea simplex</i>	(Rai and Misra, 2010)
23	Sarlahi	<i>Oscillatoria mougeotii, Spirulina major, Spirulina subsalsa</i>	(Sahay et al., 1993)
24	Sindupalchowk	<i>Oscillatoria granulate, Oscillatoria jasorvensis, Oscillatoria subbrevis, Trochiscia species, Oscillatoria limosa</i>	(Yoshimura et al., 1997; 2006)
25	Solukhumbu	<i>Aphanocapsa muscicola, Phanothece castagnei, Aphanothece nidulans, Chroococcus turicensis, Chroococcus varius, Dactylococcopsis epiphyticum, Gloeocapsopsis ferruginea, Lemmermanniella, Mantellum himalayense. Chlorogloea microcystoides, Chamaesiphon jaoi, Chamaesiphon minimus, Chamaesiphon palssahiae, Clastidium nepalense, Clastidium sicyoideum, Xenococcus luteo-violaceus, Schizothrix flammea, Schizothrix radius-solis, Anabaena laxa, Cyndrospermum breve, Cyndrospermum indicum, Coleodesmium sagarmathae, Fortiea species, Microchaete aequalis, Entophysalis rubra Want, Chroococcus tenax, Eucapsis himalayensis, Chroococcus limneticus, Stigonema mamillosum, Chlorogloea simplex, Woronichinia kuselae, Chroococcus minutus</i>	(Komarek and Watanabe, 1998; Watanabe and Komarek, 1994; Komarek and Watanabe, 1990; Subbha and Suxena, 1979)
26	Sunsari	<i>Aphanocapsa biformis, Chroococcus hangirgi, Microcystis incerta, Myxosarcina spectabilis, Arthrospira gomontiana, Oscillatoria acuta, Oscillatoria chalybea, Oscillatoria hamelii, Oscillatoria ornate, Oscillatoria vizagapatensis, Phormidium molle, Phormidium rubroterricola, Spirulina subtilissima, Symploca flotowiana, Anabaena naviculoides, Cyndrospermum stagnale, Scytonema javanicum, Gloeotrichia raciborskii, Rivularia aquatic, Microcoleus chthonoplastes Synechocystis aquatilis, Oscillatoria sancta, Spirulina gigantean, Oscillatoria chlorine, Spirulina meneghiniana, Oscillatoria splendida, Anabaena iyengarii, Spirulina subsalsa, Oscillatoria obscura, Woronichinia kuselae, Microcoleus lacustri, Spirulina princeps, Rivularia hangirgi, Gloeotrichia intermedia, Anabaena iyengarii</i>	(Jha and Kargupta, 2001)
27	Taplegunj	<i>Lyngbya limnetica</i>	(Hirano, 1963)
28	Tehrathum	<i>Phormidium favosum, Phormidium inundatum, Phormidium laminosum</i>	(Hirano, 1963)
29	Udayapur	<i>Aphanothece naegeli, Coelosphaerium dubium, Gloeotrichia echinulate, Anabaena volzii, Anabaena iyengarii</i>	(Rai and Misra, 2010)
30	Birjung	<i>Oscillatoria proboscidea, Anabaenopsis raciborskii</i>	(Prasad, 1996)
31	Rupendehi	<i>Nostoc spongiaeforme</i>	(Gyawali and Prasad, 1999)

2. APPLICATION OF ALGAE

2.1 Algae as bio- fertilizers

Bio fertilizers are the substances that comprises of microorganism, which when applied colonizes the rhizosphere and makes availability of the nutrients which ultimately helps in the growth and development of plant. It accelerates the function of microorganisms. Bio fertilizers are environment friendly and are capable of enhancing soil, crop growth and yield(Chatterjee et al., 2017). They are cheaper than other commercial insecticides and pesticides. Moreover biofertilizers do not pollute the environment like insecticides and pesticides. They are much cheaper and can be used by every farmer with general knowledge. They promote growth of plant by suppressing growth of other pathogens and diseases and are renewable sources(Scheler, 2011).

Due to the higher use of land, soil texture and fertility is decreasing day by day as a result yield is highly reduced. In order to maintain the soil quality and ultimately higher yield, algae plays a major role and acts as biofertilizers(Mahapatra et al., 2018).Algae like BGA have better capacity to fix atmospheric nitrogen under flooded condition than in dry land. An alga plays different roles and act as biofertilizers. Algae produces the adhesive substances which increases the porosity of the soil(Sahu, 2012).It excretes organic acid rich in phosphates. It tends to decreases the ph of the soil. After the death of algae biomass is added into the soil(Purwani et al., 2021). Weed growth is suppressed. the jelly structure in algae helps to hold the water, which can be used by the plants for long time (Roger and Reynaud, 1982).16. It liberates the growth hormones, vitamins, amino acid necessary for the plant growth(Rodríguez et al., 2006). As stable soil aggregates are responsible for the soil fertility, algae are responsible for the improvement of soil aggregates(Burns and Davies, 1986). Soil aggregates which are bounded by the BGA are found to have high water holding capacity(BAILEY et al., 1973).Carbon acts as the major sources for the soil bacteria , that helps in th colonization of rhizosphere through exopolysaccharides and exudates of Algae(Kapustka and DuBois, 1987).

Brown macroalgae like *Saccharina latissima*, *Laminaria digitata*, *Stoechospermum marginatum* , *Ascophyllum nodosum*, *Fucus vesiculosus*, *Ecklonia maxima*, contributes to increase nitrogen, potassium, phosphorous and carbohydrates in the soil(Eris et al., 1995; Ramya et al., 2015). Carbohydrates improves soil structure and soil aeration have good water holding capacity(Norrie & Keathley, 2006; Sharma et al., 2014). It contributes for naturally occurring hormones and growth regulators. It protects the plant from salt, freezing and drought conditions and enhances plant for photosynthetic activity and controls fungi, virus and bacteria (Craigie, 2011; Khan et al., 2011). Red Macroalgae like *Phymatolithon calcareum* (Maerla), *Lithothamnion corallioides* (Maerla) helps in the fulfilment of trace elements to soil(McHugh, 2003; Smit, 2009). Blue green algae like *Nostoc*, *Anabaena*, *Aulosira*, *Tolypothrix*, *Nodularia*, *Cylindrospermum*, *Scytonema*, *Aphanothece*, *Calothrix*, *Anabaenopsis*, *Mastigocladus*, *Fischerella*, *Stigonema*, *Haplosiphon*, *Chlorogloeopsis* fixes 18- 45 kg N/ ha in submerged rice field and produces growth promoting substances(Chatterjee et al., 2017; DE, 1939). *Anabaena Azolla* association fixes 40-80 kg of Nitrogen/ ha and can be used as green manure because of large bio mass(Moore, 1969; Vaishampayan et al., 2001).

2.2 Algae for medicines

a. Goitre treatment

Lack of iodine in body causes swelling of the muscles near the neck region which is referred as Goitre in medical terms. Algae have high amount of iodine and hence they are used to treat Goitre. (Stein and Borden 1984).

b. Obstetrical Uses

Dried stripes of brown green algae which are popularly known as 'laminaria tent' which functions as cervical dilators during obstetrical and gynecological practices (Newton 1972; Hale & Pion 1972; Feochari 1979).It is used for birth control (Manabe et al., 1982).

c. Vermifuge activity

Vermifuge is a medicine that is used to kill parasites without causing damages to the host. Red algae secretes kainic acid which helps to block neuromuscular in worm (Michanek 1979). Domic acid produced by *Digenea simplex* and *Chondria armata* behave as anthelmintic agent (Higa and Kuniyoshi 2000). Secretions from macro algae such as kainoids, terpenoids are insecticidal (Maeda et al., 1984).

d. Thalassotherapy

For the treatment of osteoporosis and rheumatism, the paste of macroalgae is used. In this method, macroalgae are crushed down into small pieces and paste is made, this paste is colled and applied over the persons body followed by warming with the help of infrared radiation (Barsanti and Gualtieri, 2006).

e. Biocosmetics

Microalgae such as *Spirulina* (Arthrospira) and *Chlorella* are used for the production of cosmetic items. Such obtained cosmetic items would work for face, skin, hands, hair and acts as refreshing items, emollient creams, anti-aging cream, and anti-irritant etc(Stolz and Obermayer 2005).

f. Antioxidant activity

Due to the excessive emotional and physical stress in daily life, it leads for the production of active free radicles. These free radicles affects the metabolism process and leads for the destruction of DNA (Barry and Aruoma, 1991), and protein synthesis.(Martin and Dean, 1991). Because of this it enhances aging process and give rise to different diseases like cancer, cardiovascular, chronic inflammation and ageing processes(Barry and Gutteridge, 1989).Algae contains antioxidants like β -carotene, astaxanthin and canthaxanthin (Borowitzka, 1988a; Lee and Zhang, 1999), polyphenols, vitamins and phycobiliproteins(Plaza et al., 2008), which helps to reduce the carcinogenesis by scavenging free radicals and active oxygen. Beta carotene present in the algae helps to prevent from leukoplakia (Sithranga Boopathy and Kathiresan, 2010).

g. Antibiotic activity

Algae maily in macroalgae have large number of chemical compounds that works as antibiotic activities (Stein and Borden, 1984; Lincoln, et al., 1991; Smit, 2004; Chanda, et al., 2010). Some green algae and microalgae produces acrylic acid from *Phaeocystis pouchettii* (Fenicol, 1975; Hashimoto, 1979) and *Chlorellin* from *Chlorella* also behaves as antibiotics. Methanol extracts from *Cladophora* helps to control *Staphylococcus aureus* and *Bacillus mycoides* activity (Demina and Mal'dov, 1981)

h. Anticardiovascular disease activity

High cholestleol levels leads to the blood pressure and ultimately causes cardiovascular diseases. The polysaccharides present in the algae like alginates carrageenans (Kiriyaama, et al., 1968), funorans, fucoidans (Lamela, et al., 1989), laminarans, porphyrans, ulvans produces hypocholesterolemic and hypolipidemic activity and reduces cholestleol level in the body(Panlasigui, et al., 2003). Systolic pressure in the body is reduced by algal polysaccharides and this activity leads to reduce the occurrence of cardiovascular diseases (Renn, et al., 1994).

i Anticoagulant and antithrombic activity

Anticoagulant and antithrombic are the terms that prevents the blood from clotting. Algae that contain sulfated polysachharides, fucoidins acts as anticoagulants and antithrombic. Fucoid helps in the synthesis of blood by regulating heparin cofactor II or antithrombin III (Church, et al., 1989; Collic, et al., 1991; Matou, et al., 2002). Major anticoagulant substances like sulfated fucans are produced from *Ascophyllum nodosum* and *Fucus vesiculosus* (Smit, 2004).

j. Anticancer and antitumor activity

the debromoaphysiatoxin obtained from the BGA showed against Lymphotic mouse leukemia (Mynderse, et al., 1977). Kahalalide F taken from *Bryopsis*(Hamann and Scheuer, 1993; Hamann, et al., 1996)., and extracts from *Chondria atropurpurea*(Palermo, et al., 1992), *Bifurcaria bifurcata*(Valls, et al., 1995), *Sargassum sp.*, *Laminaria sp.*(Yamamoto, et al., 1974, 1981) have anticancer and antitumor property which have ability to control lung colon, prostrate, nasopharyngeal cancer(Scheuer, et al., 2000).

k. Antiviral activity

Dextran sulfate, heparin, sulfated galactan, sulfated xylomannan, pentosan polysulfate from the algae found effective in controlling HIV agents (Witvrouw and De Clercq, 1997). Moreover such algae produces the enzymes that controls respiratory syncytial virus (RSV), human cytomegalovirus (HCMV), herpes simplex virus (HSV) and influenza virus (Damonte et al. 1994),*Schizymenia pacifica*, *Gracilaria corticata*, *Aghardhiella tenera*, *Nothogenia fastigata* are the important enzymes secreated by algae that mainly works as antiviral activity.

I. Anti-inflammatory activity

living tissues suffers from inflammation and it leads for the death of cell and occurrence of several diseases (Jaswir and Monsur, 2011). Macroalgae grows naturally in aquatic environment and have capacity to tolerate several stress condition and can trigger immunological responses in man (Pompani, 2001; Ehrlich 2010). Sulfated polysaccharides mainly alkaloids (Caulerpin I, II, III) obtained from red and green algae, fucoidins obtained from brown algae, and fucoxanthin and astaxanthin (carotenoids), Vidalols A and B and Phaeophytin A are anti-inflammatory compounds (Bhakuni and Rawat, 2005; Jaswir and Monsur, 2011).

2.3 Algae as food

Algae are the primary producers of ecosystem which contains chlorophyll and can prepare its food by itself. Algae are considered very important source of food to quench thirst and hunger since ancient time. During world wars, people eat *Chondrus crispus* to get energy. In Germany breads made from ground algae were prepared. *Pyropia columbina* have greater thirst-quenching tendency (Schwimmer and Schwimmer, 1955). The food obtained from the algae could be beneficial as well as harmful to the human beings and animals. The food which contains elements like cadmium or fucotoxins are harmful, which aims to protect algae from herbivores and pathogens. (Pooja, 2014) Algae are rich in protein, vitamins, minerals, amino acids, dietary fibres, lipids, polysaccharides and fatty acids. As algae do not have any circulation system, they have higher energy as compared to the higher plants. The different types of nutrients found in algae which are necessary for the animals and humans are mentioned below-

a. Polysaccharides

Polysaccharides are the chemical substance whose molecules contain bounded carbon atoms. *Odontella*, *Aphanizomenon*, *Haematococcus*, *Arthrospira*, (*Spirulina*), *Chlorella*, *Porphyridium*, *Scenedesmus*, *Dunaliella* are the algae containing polysaccharides. Green algae contains sulphuric acid polysaccharide and sulfated galactans, brown algae (Phaeophyceae) contains laminarin, fucoidan and alginic acid and red algae (Rhodophyceae) contains floridean, carrageenan and porphyran. Among the different types of algae *Ulva* species contain high quantity of polysaccharides (Pal et al., 2014). Water-soluble polysaccharides shows hypoglycemic and hypocholesterolemic effects and water-insoluble polysaccharides shows laxative effects (Burtin, 2003). Polysaccharides help to maintain sugar level by removing or adding glucose in the human body.

b) Dietary fibres

These are the substances which are indigestible obtained from the plants. Algae produces alginic acid, agars, furanans, laminarin and porphyran, cellulose, mannans and xylan as dietary fibres. These fibres help to regulate the metabolism process in the body, making it healthy. Dietary fiber content found in some algae is as shown in following tables (Mabeau and Fleurence, 1993).

Table 3: Dietary content of algae	
Algae	Dietary fibre %
Phaeophyceae	
<i>Himanthalia elongata</i>	32.70
<i>Sargassum fusiforme</i> syn. <i>Hizikia fusiforme</i>	49.20
<i>Undaria pinnatifida</i>	35.30
<i>Laminaria digitata</i> <i>Undaria</i>	38
Chlorophyceae	
<i>Ulva lactuca</i>	38
Rhodophyceae	
<i>Pyropia tenera</i>	34.80

c) Protein and amino acids

Algae contains different types of proteins and amino acids that differ from each other. Red and green algae have comparatively high amount of proteins as compared to brown algae. *Ulva* species contains 15-20 % of proteins. Amino acids like alanine, aminobutyric acid, citrulline, hydroxyl proline, ornithine and taurine as structural elements, glutamic acid, and aspartic acid are present in all algae (Wells et al., 2017). Algae can also

produce amino acids like leucine, methionine, tryptophan, lysine, threonine, and valine which human body cannot synthesize (Mabeau and Fleurence, 1993). Protein and amino acids help for the formation of new cells in the body.

d) Lipids and fatty acids

Algae contain the lipids and fats which are required for the body. Alpha linolenic acid in green algae, eicosapentaenoic acid in brown algae and docosahexaenoic acid in red algae are present (Kolanjinathan et al., 2014). All the fats and lipids while consuming by human help to regulate human physiology and metabolism. *Porphyridium cruentum*, *Nitzschia laevis*, *Phaeodactylum tricornutum* contains eicosapentaenoic acid whereas docosahexaenoic acid is found in *Cryptocodinium cohnii*, *Schizochytrium mangrovei* (Martins et al., 2013). Moreover green algae contain cholesterol, β -sitosterol and methylene cholesterol and Desmosterol, cholesterol, chalinasterol, sitosterol, and fucosterol in red algae that help to regulate blood in the body.

e) Minerals

For the proper functioning of the body, human body requires different types of minerals such as iron, iodine, phosphorous, potassium, calcium etc. This type of needs can be fulfilled while consuming macroalgae (Kolanjinathan et al., 2014). Genus *Porphyra* contains high iron, *Pseudofallacia tenera* gives high manganese, and *Seaweeds*, *Laminaria* and *Saccharina* species are nutritional sources for iodine.

f) Vitamins

Vitamins are responsible for the growth and development of the body. Deficiency of vitamins leads for the different type of diseases related to bones, eye, skin, etc. Algae such as *Palmaria palmata* and *Porphyra tenera* provide vitamins A, B1, B2 and B12. *Gracilaria chilensis* and *Codium fragile* provides pro-vitamin A (β -carotene). *Chlorella stigmatophora*, *Dunaliella tertiolecta*, *Isochrysis galbana* and *Tetraselmis suecica* possess vitamins A, B, E whereas *Ulva* and *Pyropia* sp. contains relatively less quantity of vitamin B12 (Wells et al., 2017). Brown algae contain vitamin E comparatively greater than green and red seaweeds.

2.4 Algae for biofuels

Algae are a unicellular organism that grows in moist environments. They are also considered as fastest growing organisms. Like other higher plants the process of food production of algae is similar, they utilize water, sunlight, CO₂ and other nutrients to liberate huge amount of energy and oxygen to the atmosphere. Algae can produce the renewable raw materials required for the biofuels. The oil extracted from the algae can be used directly as a biofuel including renewable diesel, jet fuels etc. The renewable diesel and jet fuels can be used directly in the engine that replaces petroleum, without causing any damages to the engines. If fuels can be obtained/produced in large quantities in near future, algae fuel can replace petroleum completely. Pyrolysis oil can be obtained from the algae that help in heat and power consumption.

Algae as biofuel have several advantages. Algae are the renewable sources of energy that can be grown everywhere using saltwater, brackish water and it does not need any special care for its growth. If we have proper amount of sunlight available we don't have any difficulty in growing algae. As it is a small plant having less metabolism activities, as a result less amount of energy is lost for its activities and huge amount of it is stored. This stored form of energy can be used for biofuels. Research shows that biofuels obtained from the algae are more than that of terrestrial plants. As carbohydrate is needed in large amount for human and animals, algae can utilize large amount of atmospheric carbon and convert into organic carbon. If we use algal biofuel then our net consumption of biofuel is nearly zero because algae are highly renewable sources of energy and can utilize it without causing any stress to environment. Process for the making algae as biofuels is similar to the purification of current biofuels. That means, we can use algal biofuel to meet the situation of other biofuels which are non-renewable in nature. Algae can be double in population within 24 hrs only so there won't be any scarcity of raw materials as biofuels. Moreover algae can be used for the production of numerous products like fertilizers, soaps etc, which requires the hydrocarbons similar to the petroleum products. Everything has both positive and negative sides. Algae too have constraints also for making algae as biofuels. Algae require large quantity of water for its proper growth and development. If less amount of water is available then, the small water temperature rises fast and which hinders the growth of algae. Algae require fertilizers like phosphorus for its proper growth. If we want to grow algae in artificial environment starting cost of production would be very large amount. We have to construct a place which holds water in high quality which is not possible with a small

investment. Algae products can be harvested daily, but you cannot utilize this product a biofuels immediately; we should follow some steps like storing in desire conditions, crushing, purifications etc for a biofuels. Although algae grows in different climates but it cannot grow in extreme climates. Cultivation of single crops for a long time causes the emergence of pest and insects. As many types of algae are found, while processing the biofuels, it creates variability in quality.

2.5 Bioremediation Of Organic Pollution

Bioremediation is a process by which substances which degrade the natural environment are break down and reduces their effects by microorganism. Algae being easily available help to fix the harmful wastage from industry, oil effulents, organic and inorganic pollutants. Different types of pollutants are described along with the potential algal species involved in their degradation.

Table 4: Pollutants absorbed by algae

Pollutant	Algae	References
Cu ²⁺	<i>Anabaena cylindrica Asterionella formosa Chlorella fusca Chlorella miniata Chlorella spp</i>	(Çetinkaya Dönmez et al., 1999; Tien et al., 2005)
Fe ³⁺	<i>Chlorella vulgaris Microcystis sp</i>	(Romera et al., 2006; S. Singh et al., 1998)
Hg ²⁺	<i>Calothrix parietina TISTR 8093 Chlamydomonas reinhardtii Cyclotella cryptica Pseudochlorococcum typicum Spirogyra hyalina</i>	(Bayramoğlu et al., 2006; Schmitt et al., 2001)
Ni ²⁺	<i>Arthrospira (Spirulina) platensis Chlamydomonas reinhardtii Chlorella miniata Chlorella spp. Chlorella vulgaris Chlorella vulgaris Chlorella vulgaris</i>	(Ferreira et al., 2011)
Pb ²⁺	<i>Arthrospira (Spirulina) platensis Chlamydomonas reinhardtii Chlorella vulgaris BCC 15 Oscillatoria laete-virens Pseudochlorococcum typicum Spirogyra insignis Spirulina (Arthrospira) platensis</i>	(Romera et al., 2007)
Zn ²⁺	<i>Aulosira fertilissima Desmodesmus pleiomorphus Isochrysis galbana Planothidium lanceolatum Spirulina spp. Stigeoclonium tenue Scenedesmus subspicatus</i>	(Singh et al., 2007)
Cd ²⁺	<i>Chaetoceros calcitrans Chlamydomonas reinhardtii Chlamydomonas reinhardtii Desmodesmus pleiomorphus (ACOI 561) Isochrysis galbana</i>	(Monteiro et al., 2010; Tüzün et al., 2005)
Cr ⁵⁺	<i>Spirulina sp. Chlamydomonas reinhardtii Chlamydomonas reinhardtii Spirulina Spirulina spp. Ulothrix tenuissima</i>	(Ming-Ju Chen, Kreuter, 1996)
Cr ₂ O ₇ ²⁻	<i>Chlorella spp. Spirulina sp.(HD-104)</i>	(Doshi et al., 2008)
Cd	<i>Cadmium(Cd) Spirogyra sps. And Oscillatoria sps.</i>	(Brahmbhatt et al, 2012)
Co, Pb,Hg	<i>Spirogyra hyaline</i>	(Shanab et al., 2012)
No ₃	<i>Chlorella vulgaris, Chlorella minutissima and Scenedesmus quadricauda</i>	(Kshirsagar, 2013)
Textile waste	<i>Chlorella vulgaris</i>	(Kassas and Mohamed, 2014)

3. FUTURE PROSPECTUS OF ALGAE

Population of the world is increasing day by day. Most of the cultivable lands are occupied by the buildings, in such conditions there can be scarcity of land for growing food. At present also we can see no cultivable land left in city areas also. Due to the increasing population scenario in near future all the areas of the world would be covered by either building or water. Due to this world could be suffered from scarcity of food. As algae can be grown in water resources, algae which contain nutritional diet for the human could fulfill this scarcity condition. Moreover algae can be used for the production of ethanol. In this context, ethanol has wide market all over the world. Algae could be the best transport fuel in near future, continuous research activities have been tested. Technology related for transportation keeps on changing, none of the technology could have dominant in the world for more than 2-3 decays. Non-renewable fuels like coal, petrol should be replaced with renewable sources of energy which are easily available in all environment conditions. Such type of fuel obtained from algae would be more effective in both cost and quality. Due to the different type of technology and works, human people may suffer from different type of diseases and can become unhealthy. Algae species can be used for the medicines. With the wider research activities more number of diseases could be identified that can be treated with algae species without any side effects. During the production of one product, industry requires waters mainly for cleaning purpose and this water is drainage into river, lakes etc. Such type of water may contain harmful chemical substances and may detriote the original quality of water. For the maintenance of water into natural form, algae could play a vital role. Due to the algae cultivation in high quantity, such type of waste water could be purified. Through the commercialization of algal cultivation, it leads for the protection of environment and minimizes risk factors. During breeding process, if we are able to insert genetic engineering and for the identification of superior quality algae then high lipid and rapid growing algae can be obtained. Algae helps for the combustion and fermentation of dead organisms, which inhibits bad, foul smell from them and gives pleasant smell. Through this environment could be natural all time. As during the consumption of every products, some byproducts are thrown out like engines emit smokes, if such fuels are made up of algae products than it won't harms environment much and contributes for sustainable

development. Establishing the algae based industries helps to provide employment opportunities to many youths. Developing algae based industries leads for the stable and overall development in economic fields in the country.

4. CONCLUSION

Algae which covers the surface of aquatic region and terrestrial region, which is distributed all over the country, can provide beneficial effect to the citizens if it can be manipulated in organized way. Proper use of modern instrument for the processing of algae can give higher return too. Algal products are beneficial, qualitative and quantative. Beneficial in a sense that for products which are obtained from the algae have some economic value. Quantative in a sense that algae are found in adverse climate, as a result algae as raw material never be diminished. Moreover algae grows faster. Algae related products have high amount of nutrient content as compared to other related products. Algae grows in unused area, due to which unused area can also give some profit, if used properly. As petroleum and diseal products which are major source of biofuel in present context, which needs a lot of effort to obtain such products. If the techniques which can process biofuel from the algae can be used, then not only it reduces the burden but also it will be qualitative, and can be easily available, which doesnot need any excavation as now. Consumption of algae in proper amount provides balance diet for the people and protects from diseases. Ultimately it helps in the purification of environment. Algae consume harmful substances secreted from different types of industry; absorb the carbondioxide and releases oxygen.

Alage are the major sources for eutrophication. During rainy seasons fertilizers like nitrogen, phosphorous are eroded from the fields, and finally reaches to the river, lakes, and oceans. Algae get such fertilizers and they grow more rapidly, they starts to absorb the oxygen found in the water and reduces the availability for aquatic organisms. As a result, aquatic organisms are hampered. Due to the presence of algae, natural environment is distrubed. Many varieties of algae are found, but in accordance with the diversity of algae very negligible research activities are performed as a result only few importances have been identified. Presence of algae makes the way slippery as a result people may slip down

and can get hurt. Algae grown in the trees, rocks can weaken the trees, rocks and mineralization may proceed in rocks. Sometimes algae may hinder the growth of main crops like rice. Inhaled of algae in high quantity can may harm health of people.

Algae have many importance rather than demerits. International as well as national organizations should focus for the research activities. Biologist working on such field should be given priority make necessary arrangement for working. They should mainly focus for algae based products and think for sustainable development. Unemployed people can be given some training for cultivation of algae; hence they can utilize their leisure time and can generate some incomes. Government should focus on the algae based industry and cultivated algae can be processed for future use. Loans for algal industry should be made interest free or reduced as far as possible from the financial sector. Technical manpower having practical knowledge should be mobilized in different remote areas, where there is high possibility of emergence of algae, and they should provide practical knowledge to the residential people regarding uses, consumption, processing, post-harvest preservation etc. If country can progress in algae industry than it would contribute a lot in economic growth and development of nation.

REFERENCES

- Ayodhya, D., Kshirsagar, 2013. Bioremediation of waste water by using microalgae: An experimental Study. *Int. J. LifeSci. Bt. and Pharma Res.* Vol. 2, No. 3.
- BAILEY, D., Mazurak, A. p., and Rosowski, J. R., 1973. j.0022-3646.1973.00099.x.pdf.
- Baral, S.R., D.K. Mishra and H.D. Kumar 1988. In situ nitrogen fixation rates in ten rice fields of Kathmandu valley, Nepal. In *Biofertilizers: Potentialities and Problems* (Eds. S.P. Sen and P. Palit). Naya Prakash, Calcutta, India. Pp. 103-107.
- Barry, H., and Gutteridge, J. M. C., 1989. In: *Free Radicals in biology and medicine*. H. Barry, ed. Oxford, Clarendon Press, Pp. 254-255.
- Barsanti, L., and Gualtieri, P., 2006. *Algae: Anatomy, Biochemistry, and Biotechnology*. Taylor and Francis. Boca Raton, USA. Pp. 301.
- Bayramoğlu, G., Tuzun, I., Celik, G., Yilmaz, M., and Arica, M. Y., 2006. Biosorption of mercury(II), cadmium(II) and lead(II) ions from aqueous system by microalgae *Chlamydomonas reinhardtii* immobilized in alginate beads. *International Journal of Mineral Processing*, 81(1), Pp. 35-43.
- Bhakuni, D. S., and Rawat, D. S., 2005. *Bioactive Marine Natural Products*. Springer, New Delhi, India.
- Black, D. L., and McGinnis, J., 2006. Abstracts. *Arabic Sciences and Philosophy*, 16(1), Pp. 3-9.
- Borowiczka, M. A., 1988. Vitamins and fine chemicals from microalgae. In: *Micro- algal Biotechnology* M. A. Borowiczka, and L. J. Borowiczka, eds., Cambridge University Press, Cambridge, Pp. 153-196.
- Brahmbhatt N.H., Patel V. Rinku and Jasrai R.T. 2012. Bioremediation Potential of *Spirogyra* sps. and *Oscillatoria* sps. for Cadmium. *Asian Journal of Biochemical and Pharmaceutical Research*. Vol 2. Issue 2.
- Burns, R. G., and Davies, J. A., 1986. The microbiology of soil structure. *Biological Agriculture and Horticulture*, 3(2-3), 95-113. Burtin, P. (2003). *Issn : 1579-4377 Nutritional Value of Seaweeds. Quality*, 2(4), 498-503
- Çetinkaya Dönmez, G., Aksu, Z., Öztürk, A., & Kutsal, T. (1999). A comparative study on heavy metal biosorption characteristics of some algae. *Process Biochemistry*, 34(9), 885-892
- Chanda, S., Dave, R., Kaneria, M. and Nagani, K. 2010. Seaweeds: A novel, untapped source of drugs from sea to combat infectious diseases. In: *Current Research, Technology and education topics in Microbiology and Microbial Biotechnology*. A. Méndez-Vilas, ed., FORMATEX, Pp.473-480.
- Chatterjee, A., Singh, S., Agrawal, C., Yadav, S., Rai, R., & Rai, L. C. (2017). Role of Algae as a Biofertilizer. *Algal Green Chemistry: Recent Progress in Biotechnology*, Pp. 189-200.
- Church F.C, Meade J. B., Treanor, R. E. and Whinna, H. C. 1989. Antithrombin activity of fucoidan. *J. biol. Chem.*, 264: Pp. 3618-3623.
- Collic, S., Fischer, A. M., Tapon-Brethaudiere, J., Boisson C, Durand, P. and Jozefonvicz, J. 1991. Anticoagulant properties of a fucoidan fraction. *Thromb. Res.*, 64: Pp. 143-154.
- Craigie, J. S. (2011). Seaweed extract stimuli in plant science and agriculture. *Journal of Applied Phycology*, 23(3), 371-393
- D. Sahu, I. P. and B. R. (2012). *Cyanobacteria - As Potential Biofertilizer*. 1, Pp. 20-26.
- Damonte, E., Neyts, J., Pujol, C. A., Snoeck, R., Andrei, G., Ikeda, S., Witvrouw, M., Reyman, D., Haines, H., Matulewicz, M. C., Cerezo, A., Coto, C. and De Clercq, E. 1994. Antiviral activity of a sulphated polysaccharide from the red seaweed *Nothogenia fastigata*. *Biochem. Pharmacol.*, 47: Pp. 2187-2192.
- Das, S.N. and B.N. Verma 1996. Algal flora of Chitwan and Nawalparasi districts of Nepal. *Phykos* 35(1-2): Pp. 119-127.
- DE, P. . (1939). The role of blue-green algae in nitrogen fixation in rice-fields. *Proceedings of the Royal Society of London. Series B - Biological Sciences*, 127(846), Pp. 121-139.
- Demina, N. S. and Mal'dov, D. G. 1981. Antibiotic properties of *Cladophora* sp. (Chlorophyceae) algae. *Izv. Akad. Nauk SSSR Ser. Biol.*, 1981: Pp. 468-470.
- Doshi, H., Seth, C., Ray, A., Kothari, I. L., 2008. Bioaccumulation of heavy metals by green algae. *Current Microbiology*, 56(3), Pp. 246-255.
- Ehrlich, H. 2010. *Biological Materials of Marine Origin invertebrate*. Volume 1 Springer Dortrecht Heidelberg London New York, Pp. 1-5.
- Elnabris. (2000). 3. Taxonomy and classification of Algae. Pp. 31-65.
- Eris, A., Sivritepe, H. Ö., Sivritepe, N., 1995. the Effect of Seaweed (*Ascophyllum nodosum*) Extract on Yield and Quality Criteria in Peppers. In *Acta Horticulturae* (Issue 412, Pp. 185-192).
- Evans Robers Jr · Denise G. Anderson · Martha T. Nester Eugene W. Nester. (2007). *Microbiology A human perspective*. 2008.
- Fall, B., Plantae, K., Kingdom, P., Protista, K., Bacteria, D., 2010. *Concepts of Botany. Algae*. 2007, Pp. 1-21.
- Fenicol, W., 1975. Halogenation in the Rhodophyta. A review. *J. Phyco.* 11: Pp. 245-259.
- Ferreira, L. S., Rodrigues, M. S., de Carvalho, J. C. M., Lodi, A., Finocchio, E., Perego, P., Converti, A., 2011. Adsorption of Ni²⁺, Zn²⁺ and Pb²⁺ onto dry biomass of *Arthrospira* (Spirulina) platensis and *Chlorella vulgaris*. I. Single metal systems. *Chemical Engineering Journal*, 173(2), Pp. 326-333.
- Francisco, C., Banaigs, B., Valls, R., and Codomier, L. 1985. Mediterranean a, a novel rearranged diterpenoid-hydroquinone from the marine alga *Cystoseira mediterranea*. *Tetrahedron Lett.*, 26: 2629-2632.
- Gyawali, Y.P. and Prasad, B.N., 1999. Survey of some potential blue-green algae from the rice fields of Butwal area and estimation of nitrogen content in *Nostoc* sp. In: *Proc. III Natn. Conf. on Sci. and Tech.*, March 8-11, 1999, RONAST, Kathmandu, Nepal. Pp. 89.
- Habib, I. 1997. Algal flora from Mahendranagar, Nepal. *J. Econ. and Taxon. Bot. (India)* 21(1): Pp. 19-26.
- Hala Yassin El- Kassas and Laila Abdelfattah Mohamed .2014. *The Egyptian Journal of Aquatic Research*. Vol. 40(3): Pp. 301-308.
- Hale, R.W. and Pion, R. J. 1972. *Laminaria: an underutilized clinical adjunct*. *Clin. Obstet. Gynecol.* 15: Pp. 829-850.
- Hamann, M. T., Scheuer, P. J., 1993. Kahalalide F: A bioactive depsipeptide from sacoglossan mollusk, *Elysia rufescens* and the green alga *Bryopsis* sp. *J. Am. Chem. Soc.*, 115: Pp. 5825-5826.
- Hamann, M. T., Otto, C. S., Scheuer, P.J. and Dunbar, D. C., 1996. Kahalalides: Bioactive peptides from a marine mollusk *Elysia rufescens* and its algal diet *Bryopsis* sp. *J. Org. Chem.*, 61: Pp. 6594-6600.

- Hasan, M. R., Chakrabarti, R., 2009. Algae FAO. Use of Algae and Aquatic Macrophytes as Feed in Small-Scale Aquaculture, Pp. 3–8.
- Hashimoto, Y., 1979. Marine toxins and other bioactive marine metabolites. Japan Scientific Society Press, Tokyo, 369 pp.
- Hickel, B. 1973. Limnological investigations in lakes of Pokhara valley, Nepal. *Int. Rev. ges Hydrobiol.* 58(5): Pp. 659-672
- Higa, T., and Kuniyoshi, M., 2000. Toxins associated with medicinal and edible seaweeds. *J. Toxicol. Tox. Rev.*, 19: Pp. 119–137.
- Hirano, M., 1955. Fresh water algae. In *Fauna and flora of Nepal Himalaya* (Ed. H. Kihara). Fauna and Flora Research Society, Kyoto University, Kyoto, Japan. Pp. 5-42.
- Hirano, M., 1963. Fresh water algae from the Nepal Himalaya, collected by a member of the Japanese Climbing Expedition. *Contr. Biol. Lab., Kyoto Univ., Japan* 16: Pp. 1-23.
- Hirano, M., 1969. Fresh water algae from Langtang Himal, Nepal Himalaya. *Contr. Biol. Lab., Kyoto Univ., Japan.* 22: Pp. 1-42.
- Jaswir, I., and Monsur, H. A., 2011. Anti-inflammatory compounds of macroalgae origin: a review. *J. Med. Pl. Res.*, 5 (33): Pp. 7146-7154.
- Jha, S., and Kargupta, A.N., 2001. Cyanobacterial flora of Eastern Koshi basin, Nepal. *Ecoprint* 8(1): Pp. 37-43.
- Joshi, A.R., 1977. Some myxophyceae of Kathmandu valley, Nepal: *Oscillatoria*. *J. Nat. Hist. Mus.* 1(1): Pp. 89-92.
- Joshi, A.R., 1979. Contribution to our knowledge on Myxophyceae of Nepal. *J. Nat. Hist. Mus. (Nepal)* 3(2): Pp. 35-41.
- Kapustka, L. A., DuBois, J. D., 1987. Dinitrogen Fixation By Cyanobacteria and Associative Rhizosphere Bacteria in the Arapaho Prairie in the Sand Hills of Nebraska. *American Journal of Botany*, 74(1), Pp. 107–113
- Khan, W., Hiltz, D., Critchley, A. T., Prithiviraj, B., 2011. Bioassay to detect *Ascophyllum nodosum* extract-induced cytokinin-like activity in *Arabidopsis thaliana*. *Journal of Applied Phycology*, 23(3), Pp. 409–414
- Kiriya, S., Okazaki, Y. and Yoshida, A., 1968. Hypocholesterolemic effect of polysaccharides and polysaccharide foodstuffs in cholesterol fed rats. *J. Nutr.*, 97: Pp. 382–388.
- Kolanjinathan, K., Ganesh, P., Saranraj, P., 2014. Pharmacological Importance of Seaweeds: A Review. *World Journal of Fish and Marine Sciences*, 6(1), Pp. 1–15.
- Komarek, J., and Lukavsky, J., 1988. *Arthronema*, a new cyanophyte genus from Afro-Asian deserts. *Arch. Hydrobiol. Suppl. Algal. Stud.* 80: Pp. 249-267.
- Komarek, J., and Watanabe, M., 1998. Contribution to the attached Cyanoprokaryotes from submerged biotopes in Sagarmatha National Park (Eastern Nepal). *Bull. Natn. Sci. Mus., Ser. B, Tokyo.* 24(4): Pp. 117-135.
- Korbitz, H., Johnson, T., Forbes, A., Brunson, M. W. C., Lutz, G., Durborow, R. M., 1994. Effects of Different Fertilizers on Algal Development. *SRAC Publication*, 466, Pp. 1–4.
- Lee, R. E., 2008. *Phycology and the algae Basic characteristics of the algae* Fourth edition Structure of the algal cell.
- Lee, Y. K., and Zhang, D. H., 1999. Production of astaxanthin by *Haematococcus*. In: *Chemicals from microalgae*. Z. Cohen, ed., Taylor and Francis, London, Pp. 173- 195.
- Lincoln, R. A., Struinski, K. and Walker, J. M., 1991. Bioactive compounds from algae. *Life Chem. Rep.*, 8: Pp. 97-183.
- Mabeau, S., Fleurence, J., 1993. Seaweed in food products: biochemical and nutritional aspects. *Trends in Food Science and Technology*, 4(4), Pp. 103–107.
- Maeda, M., Kodama, T., Tanaka, T., Ohfune, Y., Nomoto, K., Nishimura, K., and Fujita, T., 1984. Insecticidal and neuromuscular activities of domoic acid and its related compounds. *J. Pest. Sci.*, 9: Pp. 27–32
- Mahapatra, D. M., Chanakya, H. N., Joshi, N. V., Ramachandra, T. V., Murthy, G. S., 2018. Algae-Based Biofertilizers: A Biorefinery Approach. Pp. 177–196.
- Manabe, Y., Manabe, A. and Sakaguchi, M. 1982. Nelaton catheter in non-gravidas for safe and gradual cervical softening and dilation: a possible involvement of prostaglandins. *Contraception*, 25: Pp. 211-218.
- Martin, H., and Dean, M., 1991. Identification of a thiodoxin-related protein associated with plasma membranes, *Biochem. Biophys. Res. Commun.*, 175: Pp. 123-128.
- Martins, D. A., Custódio, L., Barreira, L., Pereira, H., Ben-Hamadou, R., Varela, J., Abu-Salah, K. M., 2013. Alternative sources of n-3 long-chain polyunsaturated fatty acids in marine microalgae. *Marine Drugs*, 11(7), Pp. 2259–2281
- Matou, S., Helley, D., Chabut, D., Bros, A., Fischer, A-M., 2002. Effect of fucoidan on fibroblast growth factor-2-induced angiogenesis in vitro. *Thromb. Res.*, 106: Pp. 213–221.
- McHugh, D. J., 2003. Seaweeds uses as Human Foods. In *A Guide to the Seaweed Industry* (Issue 441).
- Michanek, G., 1979. Sea weed resources for pharmaceutical uses. In: *Marine algae in Pharmaceutical Science*. H. A. Hoppe, T. Levring, and Y. Tanaka, eds., Walter de Gruyter, Berlin, Pp. 203-235.
- Ming-Ju Chen, Kreuter, J. Y.-T. K., 1996. Nanoparticles and microparticles for drug and vaccine delivery. *Journal of Anatomy*, 189 (Pt 3(Ii), Pp. 503–505.
- Monteiro, C. M., Castro, P. M. L., Malcata, F. X., 2010. Cadmium Removal by Two Strains of *Desmodesmus pleiomorphus* Cells. *Water, Air, and Soil Pollution*, 208(1–4), Pp. 17–27.
- Moore, A. W., 1969. *Azolla: Biology and agronomic significance*. The Botanical Review, 35(1), Pp. 17–34.
- Munir, N., Sharif, N., Naz, S., Farkhanda, M., 2013. Algae: A potent antioxidant source. *Sky J Microbiol Res*, 1(3), Pp. 22–31. [http://www.skyjournals.org/sjmr/pdf/2013pdf/Apr/Munir et al pdf.pdf](http://www.skyjournals.org/sjmr/pdf/2013pdf/Apr/Munir%20et%20al.pdf)
- Mynderse, J. S., Moore, R. E., Kashiwagi, M. and Norton, T. R., 1977. Antileukemia activity in *Oscillatoriaceae*: isolation of debromoaplysiatoxin from *Lyngbya*. *Science*, 196: Pp. 538-539.
- Nakanishi, M., 1986. Limnological study in Phewa, Begnas and Rupa lakes. In *Studies on distribution, adaptation and evolution of microorganisms in Nepal Himalayas* (Ed. Y. Ishida). (Second report), Ministry of Education, Science and Culture, Kyoto, Japan. Pp. 3-13.
- Newton, B. W., 1972. *Laminaria* tent: relic of the past or modern medical device. *Am. J. Obstet. Gynecol.*, 113: Pp. 442-448.
- Norrie, J., Keathley, J. P., 2006. Benefits of *Ascophyllum nodosum* marine-plant extract applications to Thompson seedless grape production. *Acta Horticulturae*, 727(November 2006), Pp. 243–247. <https://doi.org/10.17660/actahortic.2006.727.27>
- Pal, A., Kamthania, M. C., Kumar, A., 2014. Bioactive Compounds and Properties of Seaweeds—A Review. *OALib*, 01(04), Pp. 1–17. <https://doi.org/10.4236/oalib.1100752>
- Palermo, J. A., Flower, B. P., Seldes, A. M., 1992. Chondriamides A and B, new indolic metabolites from the red alga *Chondria* sp. *Tetrahedron Lett.*, 33: Pp. 3097- 3100.
- Pant, K., and Gupta, V.N.P., 1999. Blue-green algae floristics of the rice fields of Lamjung (Nepal). In: *Proc. III National Conf. on Science and Technology*, Abs. BSC 013. Royal Nepal Academy of Science and Technology (RONAST), Kathmandu, Nepal, Pp. 70.
- Plaza, M., Cifuentes, A., Ibáñez, E., 2008. In the search of new functional food ingredients from algae. *Trends in Food Science and Technology*, 19(1), Pp. 31–39.
- Pompani, S. A., 2001. The oceans and human health: the discovery and development of marine-derived drugs. *Ocenography*, 14: Pp. 78-87.
- Pooja, S., 2014. Algae used as Medicine and Food-A Short Review. *Journal of Applied Pharmaceutical Science and Research*, 6(162), Pp. 33–35.
- Prasad, R.C., Prasad, B.N., 2001. Screening of blue green algae (Cyanobacteria) and their distributional pattern in rice field of Narayani and Bagmati zones of Nepal. *J. Liv. World* 8(1): Pp. 1-12.

- Prasad, V., 1996. Blue green algae of Birganj (Nepal). Short term project. Research Division, Tribhuvan University, Nepal.
- Purwani, J., Pratiwi, E., Sipahutar, I. A., Husnain, 2021. The effect of different species of cyanobacteria on the rice yield and nitrogen use efficiency under different levels of nitrogen fertilizer on Alluvial West Java. IOP Conference Series: Earth and Environmental Science, 648(1)
- Rai, S.K., Misra, P.K., 2010. Freshwater cyanophyceae from east Nepal. Bangladesh J. of Plant Taxonomy 17(2): Pp. 121-139.
- Ramya, S. S., Vijayanand, N., Rathinavel, S., 2015. Foliar application of liquid biofertilizer of brown alga *Stoechospermum marginatum* on growth, biochemical and yield of *Solanum melongena*. International Journal of Recycling of Organic Waste in Agriculture, 4(3), Pp. 167–173
- Renn, D. W., Noda, H., Amano, H., Nishino, T., Nishizana, K., 1994a. Antihypertensive and antihyperlipidemic effects of funoran. Fisheries Sci., 60: Pp. 423–427.
- Rodríguez, A. A., Stella, A. M., Storni, M. M., Zulpa, G., Zaccaro, M. C., 2006. Effects of cyanobacterial extracellular products and gibberellic acid on salinity tolerance in *Oryza sativa* L. 4, Pp. 1–4.
- Roger, P. A., Reynaud, P., 1982. Microbiology of Tropical Soils and Plant Productivity. Microbiology of Tropical Soils and Plant Productivity, January.
- Romera, E., González, F., Ballester, A., Blázquez, M. L., Muñoz, J. A., 2006. Biosorption with algae: A statistical review. Critical Reviews in Biotechnology, 26(4), Pp. 223–235
- Romera, E., González, F., Ballester, A., Blázquez, M. L., Muñoz, J. A., 2007. Comparative study of biosorption of heavy metals using different types of algae. Bioresource Technology, 98(17), Pp. 3344–3353
- Scheler, M., 2011. Biofertilizers. 8(4), 2014.
- Scheuer, P. J., Haman, M.T. and Gravalos, D.G., 2000. Cytotoxic and antiviral compound. US Patent Number 6011010.
- Schmitt, D., Müller, A., Csögör, Z., Frimmel, F. H., Posten, C., 2001. The adsorption kinetics of metal ions onto different microalgae and siliceous earth. Water Research, 35(3), Pp. 779–785
- Schwimmer, M., Schwimmer, D., 1955. The Role of Algae and Plankton in Medicine.
- Shanab, S., Essa, A., Shalaby, E., 2012. Bioremoval capacity of three heavy metals by some microalgae species (Egyptian isolates). Plant Signaling and Behavior, 7(3).
- Sharma, H. S. S., Fleming, C., Selby, C., Rao, J. R., Martin, T., 2014. Plant biostimulants: A review on the processing of macroalgae and use of extracts for crop management to reduce abiotic and biotic stresses. Journal of Applied Phycology, 26(1), Pp. 465–490.
- Shrestha, B., Manandhar, J.D., 1983. Contribution to the algal flora of Kathmandu valley. J. Inst. Sci. Techn. T.U. (Nepal) 6: Pp. 1-6
- Singh, A., Mehta, S. K., Gaur, J. P., 2007. Removal of heavy metals from aqueous solution by common freshwater filamentous algae. World Journal of Microbiology and Biotechnology, 23(8), Pp. 1115–1120.
- Singh, S., Pradhan, S., Rai, L. C., 1998. Comparative assessment of Fe³⁺ and Cu²⁺ biosorption by field and laboratory-grown *Microcystis*. Process Biochemistry, 33(5), Pp. 495–504
- Sithranga Boopathy, N., Kathiresan, K., 2010. Anticancer drugs from marine flora: An overview. Journal of Oncology, Pp. 1–10
- Smit, A. J. 2004. Medicinal and pharmaceutical uses of seaweed natural products: a review. J. Appl. Phycol., 16: Pp. 245–262.
- Smit, A. J., 2004. Medicinal and pharmaceutical uses of seaweed natural products: a review. J. Appl. Phycol., 16: Pp. 245–262.
- Smit, A. W., 2009. What is the potential of marine alga, in combination with sewage sludge, as a composite of bio-fertilizer?
- Stein, J. R., and Borden, C. A., 1984. Causative and beneficial algae in human disease condition: a review. Phycologia, 23(4): Pp. 485–501.
- Stolz, P., and Obermayer, B., 2005. Manufacturing microalgae for skin care. Cosmetics Toiletries, 120: Pp. 99–106.
- Subba Raju, N., and Suxena, M.R., 1979. Algae and testacea of the Cho Oyu (Himalayas) Expedition-II; cyanophyta, chlorophyta, euglenophyta, chrysophyta and testacea. Hydrobiologia 67(2): Pp. 141–160.
- Tien, C. J., Sigee, D. C., White, K. N., 2005. Copper adsorption kinetics of cultured algal cells and freshwater phytoplankton with emphasis on cell surface characteristics. Journal of Applied Phycology, 17(5), Pp. 379–389
- Tüzün, I., Bayramoğlu, G., Yalçın, E., Başaran, G., Çelik, G., Arica, M. Y., 2005. Equilibrium and kinetic studies on biosorption of Hg(II), Cd(II) and Pb(II) ions onto microalgae *Chlamydomonas reinhardtii*. Journal of Environmental Management, 77(2), Pp. 85–92.
- Upadhyaya, B.N., 1979. Two new records of *Oscillatoria* for Nepal. J. Nat. Hist. Mus. (Nepal) 3(3): Pp. 74–75.
- Vaishampayan, A., Singh, R. P., D.P. H., Dey, T., Gupta, A. ., Bhan, U., Rao, A., 2001. 426.pdf.
- Valls, R., Piovetti, L., Banaigs, B., Archavlis, A. and Pellegrini, M., 1995. (S):13- hydroxygeranylgeraniol-derived furanoditerpenes from *Bifurcaria bifurcata*. Phytochemistry, 39: Pp. 145–149.
- Witvrouw, M., and De Clercq, E., 1997. Sulphated polysaccharides extracted from sea algae as potential antiviral drugs. Gen. Pharmac., 29: Pp. 497–511.
- Witvrouw, M., Desmyter, J., and De Clercq, E., 1994. Antiviral portrait series: 4. Polysulphates as inhibitors of HIV and other enveloped viruses. Antiviral Chem. Chemother., 5: Pp. 345–359.

